



TITLE:

# Analysis of interior-point-paths for sufficient linear complementarity problems(Continuous and Discrete Mathematical Optimization)

AUTHOR(S):

Stoer, J.

---

CITATION:

Stoer, J.. Analysis of interior-point-paths for sufficient linear complementarity problems(Continuous and Discrete Mathematical Optimization). 数理解析研究所講究録 1997, 981: 36-36

ISSUE DATE:

1997-03

URL:

<http://hdl.handle.net/2433/60899>

RIGHT:

# Analysis of interior-point-paths for sufficient linear complementarity problems

J. Stoer

Institut für Angewandte Mathematik und Statistik  
Universität Würzburg, Würzburg, Germany

**Abstract:** In this lecture we describe the behavior of infeasible-interior-point-paths for solving horizontal linear complementarity problems

$$(LCP) \quad Px + Qy = q, \quad (x, y) \geq 0, \quad x^T y = 0,$$

that are sufficient in the sense of Cottle, Pang and Venkateswaran (1989). These paths are defined as the solution  $(x, y)(r, \eta)$ ,  $r > 0$ ,  $\eta > 0$ , of

$$\begin{aligned} Px + Qy &= q + r\bar{q}, \quad (x, y) \geq 0, \\ x_i y_i &= r\eta_i, \quad i = 1, \dots, n, \end{aligned}$$

and they converge to a central point of the set of solutions of  $(LCP)$  as  $r \downarrow 0$ . It is shown that these paths are analytic functions of  $r$  even at  $r = 0$ , if  $(LCP)$  has a strictly complementary solution, and are analytic in  $\rho := \sqrt{r}$  at  $\rho = 0$ , if  $(LCP)$  is solvable but has no strictly complementary solutions.